

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

REGION IX

75 Hawthorne Street San Francisco, CA 94105-3901

August 1, 1994

Subject:

Region IX Preliminary Remediation Goals (PRGs) Second Half 1994

From:

Stanford J. Smucker, Ph.D. Regional Toxicologist (H-9-3) Technical Support Section

To:

PRG Table Mailing List

Please find the update to the Region IX PRG table. The table has been revised to reflect the most current EPA toxicological and risk assessment information. Updates to EPA toxicity values were obtained from IRIS through July 1994 and HEAST through March 1994. Age-adjustment factors have been added to better evaluate residential exposures to carcinogens. In addition, the soll saturation equation has been corrected, leading to approximately a ten fold higher saturation concentration in soils.

The PRG table provides useful risk-based information for Region IX risk assessors and managers. However, the table has no official status and may be in conflict with local state requirements. Four problem chemicals (cadmium, chromium, nickel, and DBCP) have been identified by Cal-EPA's Department of Toxic Substances Control. California soil values differ significantly, by a factor of four or more, for these chemicals. To address these concerns, the "Cal-modified PRG" concentrations, based on PEA (1994) guidance, are included with the federal values and should be used in California when screening a site.

In general, PRGs should be used as a predictor of single-contaminant risk estimates for a specific environmental media (e.g. soil, air, and tap water). However, multiple pollutant risks can also be estimated using PRGs (see Screening Risk below). This procedure is somewhat more complicated as it requires gathering additional information, either by downloading the table to display the hidden columns or by using the equations presented in the text for calculating additional concentration terms not provided in the print out.

A contaminant concentration that exceeds a PRG level does not, in itself, mean that there is an unacceptable health threat. However, exceedances should be evaluated further. It is recommended that the reader verify the numbers with a toxicologist because the toxicity/exposure information in the table may contain errors or default assumptions that need to be refined based on further evaluation.

If you are not currently on the PRG mailing list, but would like to be, please make the request through EPA's project manager working on your site. Or, simply download the file (PRG2ND94,ZIP) from California Regional Water Board's BBS [(510) 286-0404]. If you find an error please send me a note via fax at (415) 744-1916.

1.0 INTRODUCTION

The Region IX PRG Table combines EPA toxicity values, updated blannually, with reasonable maximum exposure (RME) factors to estimate concentrations in environmental media (e.g. soil, air, and water) that are generally agreed to be "safe" for humans. Above these levels, there may be enough concern to warrant further evaluation of risks.

PRG concentrations presented in the Tables can be used to screen pollutants in environmental media, trigger further Investigation, and provide an initial cleanup goal if applicable.

When considering PRGs as Initial cleanup goals, residential concentrations should be used for maximum beneficial uses of a property. Industrial concentrations for soil only are included in the table as an alternative goal, but industrial concentrations should not be used for screening a site. They are meant to provide the manager with an alternative preliminary goal for sites zoned heavy industrial.

Before applying PRGs as screening tools or initial cleanup goals, the user of the table should consider whether the exposure pathways at the site are fully accounted for in the PRG calculation. Region IX PRG concentrations are based on direct exposures (i.e. ingestion, dermal contact, and inhalation) for specific land-use conditions and do not consider impact to groundwater or ecological receptors. To determine the appropriateness of Region IX PRGs, the following questions should be asked:

- Are there potential ecological concerns?
- Is there potential for land use other than those covered by the PRGs (that is, residential and industrial)?
- Are there other likely human exposure pathways that were not considered in development of the PRGs (e.g. impact to groundwater, local fish consumption; raising beef, dairy, or other livestock)?
- Are there unusual site conditions (e.g. large areas of contamination, high fugitive dust levels, potential for indoor air contamination)?

If any of these four conditions exist, the PRG may need to be modified to reflect this new information. In general, PRGs are refined in the site conceptual model developed as part of a site-specific risk assessment.

DISCLAIMER

Preliminary remediation goals (PRGs) focus on dominant exposure pathways and may not consider all exposure pathways encountered at CERCLA/RCRA sites (Exhibit 1-1). PRGs do not consider impact to groundwater or address ecological concerns. PRGs are specifically not intended as a (1) stand-alone decision-making tool, (2) as a substitute for EPA guidance for preparing baseline risk assessments, or (3) a rule to determine if a waste is hazardous under RCRA.

The guidance set out in this document is not final Agency action. It is not intended, nor can it be relied upon to create any rights enforceable by any party in litigation with the United States. EPA officials may decide to follow the guidance provided herein, or act at variance with the guidance, based on an analysis of specific circumstances. The Agency also reserves the right to change this guidance at any time without public notice.

EXHIBIT 1-1 TYPICAL EXPOSURE PATHWAYS BY MEDIUM FOR RESIDENTIAL AND INDUSTRIAL LAND USES*

	EXPOSURE PATHWAYS, ASSUN	ing:
MEDIUM	RESIDENTIAL LAND USE	INDUSTRIAL LAND USE
Ground Water	Ingestion from drinking	ingestion from drinking
	inhalation of volatiles	Inhalation of volatiles
	Dermal absorption from bathing	Dermal absorption
Surface Water	Ingestion from drinking	Ingestion from drinking
	inhalation of volatiles	Inhalation of volatiles
	Dermal absorption from bathing	Dermal absorption
	Ingestion during swimming	
	Ingestion of contaminated fish	
Soil	Ingestion	Ingestion
	Inhalation of particulates	Inhalation of particulates
	Inhalation of volatiles	inhalation of volatiles
	Exposure to indoor air from soil gas	Exposure to indoor air from soli gas
	Exposure to ground water contaminated by soil leachate	Exposure to ground water contaminated by soil leachate
	Ingestion via plant uptake	Inhalation of particulates from trucks and heavy equipment
	Dermal absorption	Dermal absorption

Footnote:

^{*}Exposure pathways considered in the PRG calculations are indicated in boldface Italics.

2.0 READING THE PRG TABLE

2.1 General Considerations:

PRGs are health-based concentrations that correspond to either a 1 in a million (10°) cancer risk or a "safe" reference dose (RfD), whichever is lower. PRG concentrations based on cancer and noncancer concerns are indicated by "ca" and "nc", respectively. Cancer-causing agents may have additional non-cancer PRGs not listed in the Tables. These can be obtained by downloading file (PRG2ND94.ZIP) from California Regional Water Board's Bulletin Board System at [(510)286-0404)] or using the calculations provided below.

In general, PRG concentrations in the table are risk-based but for soil there are two important exceptions: 1) for several volatile chemicals PRGs are based on soil saturation equation ("sat") (see below), and 2) for relatively less toxic inorganic and semivolatile contaminants, a non-risk based "ceiling limit" concentration is given as 10⁻⁵mg/kg "max". PRG concentrations that are not risk-based (i.e. either "sat" or "max") should be segregated before screening multiple pollutant risks.

2.2 Toxicity Values:

EPA toxicity values, known as noncarcinogenic reference doses (RfD) and carcinogenic slope factors (SF) were obtained from IRIS through July 1994, HEAST through March 1994, and ECAO-Cincinnati. The priority among sources of toxicological constants used are as follows: (1) IRIS (indicated by "i"), (2) HEAST ("h"), (3) ECAO-Cincinnati ("e"), and (4) withdrawn from IRIS or HEAST ("x"). Note in some cases, state toxicity values may differ from the federal numbers or even be promulgated as ARARs; these and the resultant PRGs should also be considered as initial cleanup goals.

Route-to-route extapolations ("r") were frequently used when there were no toxicity values available for a given route. Oral cancer slope factors ("oSF") and reference doses ("oRfD") were used for both oral and inhaled exposures for organic compounds lacking inhalation values. Also, inhalation slope factors ("iSF") and inhalation reference doses ("iRfD") were often used for both inhaled and oral exposures for organic compounds lacking oral values.

An additional route-to-route extrapolation is the use of oral toxicity values for evaluating dermal exposures. Although route-to-route methods are a useful screening procedure, the assumptions may need to be further evaluated in a site-specific risk assessment.

2.3 Soll Factors:

Chemical-specific information for solls, volatilization factors ("VF_s") and skin absorption factors ("ABS"), are listed in the table to provide additional assumptions used to calculate soil PRGs. For volatile chemicals, the "VF_s" term was incorporated into the PRG equations to address long-term inhalation exposures. Volatile organic chemicals (VOCs) are indicated by "1" in the VOC column of the Table and are defined as those chemicals having a Henry's Law constant greater than 10⁻⁶ (atm-m³/mol) and a molecular weight less than 200 g/mole).

Chemical-specific "ABS" values are provided for arsenic, cadmium, pentachlorophenol, PCBs, and dioxin as recommended by EPA's Office of Research and Development (1994) for the evaluation of contaminant absorption through the skin. Otherwise, default skin absorption fractions are assumed to be 0.01 and 0.10, for inorganics and organics, respectively.

2.4 Risk Screening:

A suggested stepwise approach for screening sites with PRGs is as follows:

- Perform an extensive records search and compile existing data
- Identify site contaminants in the PRG Table. Record the PRG concentrations for various media and note whether PRG is based on cancer risk (indicated by "ca") or noncancer hazard (indicated by "nc"). Segregate cancer PRGs from non-cancer PRGs and exclude (but don't eliminate) non-risk based PRGs ("sat" or "max").
- For cancer risk estimates, take the site-specific concentration (maximum or 95 UCL) and divide by the PRG concentrations that are designated for cancer evaluation ("ca").
 Multiply this ratio by 10⁻⁶ to estimate chemical-specific risk. For multiple pollutants, simply add the risk for each chemical:

$$Risk=10^{-6} \cdot \left[\left(\frac{conc_x}{PRG_x} \right) + \left(\frac{conc_y}{PRG_y} \right) + \left(\frac{conc_z}{PRG_z} \right) \right]$$

• For non-cancer hazard estimates. Divide concentration term by its respective non-cancer PRG designated as "nc" and sum the ratios for multiple contaminants. [Note that carcinogens may also have an associated non-cancer PRG that is not listed in the printed copy of the table and these will also need to be obtained in order to complete the non-cancer evaluation.] The non-cancer ratio represents a hazard index (HI). A hazard index of 1 or less is generally considered safe. A ratio greater than 1 suggests further evaluation:

Hazard Index=
$$\left[\left(\frac{conc_x}{PRG_x}\right) + \left(\frac{conc_y}{PRG_y}\right) + \left(\frac{conc_z}{PRG_z}\right)\right]$$

For more information on screening site risks, the reader should contact EPA Region IX's Technical Support Section.

3.0 TECHNICAL SUPPORT DOCUMENTATION

PRGs consider direct exposure hazards to chemicals from contact with complex media, soils, air, and water. The emphasis of the PRG equations and technical discussion are aimed at developing initial goals for soils, since this is an area where few standards exist. For air and water, additional reference concentrations or standards are available for many chemicals (e.g. MCLGs and NAAQS) and consequently the discussion of these media are brief.

3.1 Volatile Chemicals in Soil and Water:

Volatile chemicals, defined as those chemicals having a Henry's Law constant greater than 10⁻⁶ (atm-m³/mol) and a molecular weight less than 200 g/mole, were screened for inhalation exposures using a volatilization factor in the PRG calculations for soil and water (RAGS Part B).

Volatilization factors for soils (VF_s) are chemical-specific and were calculated from physical-chemical information obtained from a number of sources including Superfund Exposure Assessment Manual (reference "1")(SEAM, EPA 1988), Superfund Public Health Evaluation Manual (reference "2")(EPA 1986), Subsurface Contamination Reference Guide (reference "3")(EPA 1990) and Fate and Exposure Data (reference "4")(Howard 1991) and are presented in Attachment A. In those cases where Diffusivity Coefficients (Di) were not provided in existing literature, Di's were calculated using Fuller's Method described in SEAM. A surrogate VF for contaminants in soil was required for some chemicals that lacked physico-chemical information. In these cases, a proxy chemical of similar structure was used that may over- or under-estimate the PRG for soils.

The basic principle of the VF model is applicable only if the soil contaminant concentration is at or below soil saturation. If the PRG calculated using VF_s was greater than the calculated "sat", the PRG was set equal to "sat" in accordance with Risk Assessment Guidance for Superfund - Part B (EPA, 1991).

For tap water, an upperbound volatilization constant (VF_w) is used that is based on all uses of household water (e.g showering, laundering, and dish washing). Certain assumptions were made. For example, it is assumed that the volume of water used in a residence for a family of four is 720 L/day, the volume of the dwelling is 150,000 L and the air exchange rate is 0.25 air changes/hour (Andelman in RAGS Part B). Furthermore, it is assumed that the average transfer efficiency weighted by water use is 50 percent [i.e. half of the concentration of each chemical in water will be transfered into air by all water uses. Note: the range of transfer efficiencies extends from 30% for toilets to 90% for dishwashers.

3.2 Dermal Absorption of Contaminants in Soil:

Much uncertainty surrounds the determination of hazards associated with skin contact with soils. Thus far, chemical-specific absorption values for skin have been recommended for only five chemicals by EPA's Office of Research and Development. For all other chemicals, default absorption values for inorganics and organics are assumed to be 1 and 10 percent, respectively. An additional uncertainty is the lack of toxicity values for the dermal route. For screening purposes it is assumed that dermal toxicity values can be route-to-route extrapolated from oral values, but this may not always be an appropriate assumption and should be checked.

At 10 % skin absorption, the dermal dose is estimated to equal an ingestion dose for adults, using the best estimate default values in *Dermal Exposure Assessment: Principles and Applications* (EPA 1992). At 1 % absorption, the dermal dose is estimated to be 10% of the oral dose (i.e. based on an adult

ingestion rate of 100 mg/day). Note: worker and children intake rates, 50 mg/day and 200 mg/day, respectively, yield somewhat different results.

dermal dose = ingestion dose

$$ABS = \frac{(100mg/day)}{[(0.2mg/cm^2-day)(5000cm^2)]} = 0.10$$

3.3 Chemicals Adsorbed to Airborne Particles:

Inhalation of chemicals adsorbed to respirable particles (PM₁₀) were assessed using a default particulate emission factor (PEF) equal to 4.63 x 10° m³/kg that relates the contaminant concentration in soil with the concentration of respirable particles in the air due to fugitive dust emissions from contaminated soils. The relationship is derived by Cowherd (1985) for a rapid assessment procedure applicable to a typical hazardous waste site where the surface contamination provides a relatively continuous and constant potential for emission over an extended period of time (e.g. years). This may not be an appropriate assumption for all sites.

With the possible exception of cadmium, chromium, and nickel, inhalation of airborne particles (under typical conditions) does not significantly affect the PRG for soils. For more details regarding specific parameters used in the PEF model, the reader is referred to RAGS Part B (EPA, 1991).

3.4 Exposure Factors:

Default exposure factors were obtained primarily from RAGS Supplemental Guldance Standard Default Exposure Factors (OSWER Directive, 9285.6-03) dated March 25, 1991 and supplemented with more recent information from U.S. EPA's Office of Solid Waste and Emergency Response, U.S. EPA's Office of Research and Development, and California EPA's Department of Toxic Substances Control (see Exhibit 3-1).

Because contact rates may be different for children and adults, carcinogenic risks during the first 30 years of life were calculated using age-adjusted factors. Use of age-adjusted factors are especially important for soil ingestion exposures, which are higher during childhood and decrease with age. However, for purposes of combining exposures across pathways, additional age-adjusted factors are used for Inhalation and dermal exposures. These factors approximate the integrated exposure from birth until age 30 combining contact rates, body weights, and exposure durations for two age groups small children and adults. Age-adjusted factors were obtained from RAGS PART B or developed by analogy.

(1) ingestion([mg-yr]/[kg-d]:

$$IFS_{adj} = \frac{ED_c \cdot IRS_c}{BW_c} + \frac{(ED_s - ED_c) \cdot IRS_a}{BW_a}$$

(2) skin contact([mg*yr]/[kg*d]:

$$SFS_{adj} = \frac{ED_c \cdot SL \cdot SA_c}{BW_a} + \frac{(ED_z - ED_c) \cdot SL \cdot SA_a}{BW_a}$$

(3) inhalation ([m³-yr]/[kg-d]):

$$InhF_{adj} = \frac{ED_c \cdot IRA_c}{BW_c} + \frac{(ED_z - ED_c) \cdot IRA_a}{BW_a}$$

For soils only, noncarcinogenic contaminants are evaluated in children separately from adults. No ageadjustment factor is used in this case. The focus on children is considered protective of the higher daily intake rates of soil by children and their lower body weight. For maintaining consistency, when evaluating soils, dermal and inhalation exposures are also based on childhood contact rates.

3.5 PRG Equations:

The equations used to calculate the PRGs for carcinogenic and noncarcinogenic contaminants are presented in Equations 3-1 thru 3-8. Calculations of PRGs are consistent with RAGS Part B (U.S. EPA 1991) but also consider updates to the RAGS Part B equations. Briefly, the methodology backcalculates a soil, air, or water concentration level from a target risk (for carcinogens) or hazard quotient (for noncarcinogens). The equations for soil combine across pathways for direct exposures (i.e. ingestion, skin contact, and inhalation). To evaluate route-specific contribution to the PRG concentration, the reader may want to download the PRG table from California Regional Water Board's BBS mentioned above and display the hidden columns.

To calculate PRGs for volatile chemicals in soil, a chemical-specific volatilization factor is calculated per Equation 3-9 (page 12). Because of its reliance on Henry's law, the VF model is applicable only when the contaminant concentration in soil water is at or below saturation (i.e. there is no free-phase contaminant present). This corresponds to the contaminant concentration in soil at which the adsorptive limits of the soll particles and the solubility limits of the available soil moisture have been reached. Above this point, pure liquid-phase contaminant is expected in the soil. The updated equation for deriving C_{ast} is presented in Equation 3-10.

EXHIBIT 3-1 STANDARD DEFAULT FACTORS

Symbol	Definition (units)	Default	Reference
CSFo	Cancer slope factor oral (mg/kg-d)-1	-	IRIS, HEAST, or ECAO
CSFI	Cancer slope factor inhaled (mg/kg-d)-1	-	IRIS, HEAST, or ECAO
RiDo	Reference dose oral (mg/kg-d)	_	IRIS, HEAST, or ECAO
RIDI	Reference dose inhaled (mg/kg-d)	-	IRIS, HEAST, or ECAO
TR	Target cancer siek	10-6	
THQ	Target hazard quotient	1	
BWa	Body weight, adult (kg)	70°	RAGS (Part A), EPA 1989 (EPA/540/1-89/002)
BWc	Body weight, child (kg)	15	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
AT	Averaging time - cancer (years)	70	RAGS(Part A), EPA 1989 (EPA/540/1-89/002)
SAB	25% Surface area, adult (cm²)	5000	Dermai Assessment, EPA 1992 (EPA/600/8-91/011B)
SAC	25% Surface area, child (cm²)	2000	Dermal Assessment, EPA 1992 (EPA/ 600/8-9/011B)
AF	Adherence factor (mg/cm²)	0.2	Dermal Assessment, EPA 1992 (EPA/ 600/8-9/011B)
ABS	Skin absorption (unitiess):		PPA G. I PPA (PPAG 4004)
	- organics	0.1	PEA, Cal-EPA (DTSC, 1994)
	-Inorganics	0.01	PEA, Cal-EPA (DTSC, 1994)
IRAs	Inhalation rate - adult (m²/day)	20	Exposure Factors, EPA 1991 (OSWER No. 9285.6-03)
IRAc	Inhalation rate - child (m²/day)	10	RAGS (Part A), EPA 1989 (EPA/540/1-89/002)
IRWa	Drinking water ingestion - adult (L/day	2	RAGS(Part A), EPA 1989 (EPA/540/1-89/002)
IRWc	Drinking water ingestion - child (L/day)	1	PEA, Cal-EPA (DTSC, 1994)
IRSa	Soil ingestion - adult (mg/day)	100	Exposure Factors, EPA 1991 (OSWER No. 9285.6-03)
IRSc	Soil ingestion - child (mg/day),	200	Exposure Factors, EPA 1991 (OSWER No. 9285.6-03)
IRSo	Soil ingestion - occupational (mg/day)	50	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
EFr	Exposure frequency - residential (d/y)	350	Exposure Factors, EPA 1991 (OSWER No. 9285.6-03)
EFo	Exposure frequency - occupational (d/y)	250	Exposure Factors, EPA 1991 (OSWER No. 9285.6-03)
EDr	Exposure duration - residential (years)	30	Exposure Factors, EPA 1991 (OSWER No. 9285.6-03)
EDc	Exposure duration - child (years)	6	Exposure Factors, EPA 1991 (OSWER No. 9285.6-03)
EDo	Exposure duration - occupational (years)	25	Exposure Factors , EPA 1991 (OSWER No. 9285.6-03)
	Age-adjusted factors for carcinogens:		
IFSadj	Ingestion factor, solis ([mg-yr]/[kg-d])	114	RAGS(Part B) , EPA 1991 (OSWER No. 9285.7-01B)
SFSadj	Skin contact factor, solis ([mg*yr]/[kg*d])	503	By analogy to RAGS (Part B)
InhFadj	Inhalation factor ([m²-yr]/[kg-d])	11	By analogy to RAGS (Part B)
VFw	Volatilization factor for water (unitiess)	0.5	RAGS(Part B) , EPA 1991 (OSWER No. 9285.7-01B)
PEF	Particulate emission factor (m²/kg) Se	e below	RAGS(Part B) , EPA 1991 (OSWER No. 9285.7-01B)
VFs	Volatilization factor for soil (m³/kg) Se	e below	OSWER (EPA 1993, communication from Janine Dinan)
Ceat	Soil saturation concentration (mg/kg) Se	e below	OSWER (EPA 1994, communication from Janine Dinan)

Footnote:

*Seventy years is the averaging time for carcinogens. For noncarcinogens, the averaging time is set equal to the exposure duration (AT = ED).

*Exposure duration for lifetime residents is assumed to be 30 years total. For carcinogens, exposures are combined for children (6 years) and adults (24 years).

PRG EQUATIONS

Soil Equations: For soils, equations were based on three exposure routes (ingestion, skin contact, and inhalation).

Equation 3-1: Direct Exposures to Carcinogenic Contaminants in Residential Soil

$$C(mg/kg) = \frac{TR\cdot AT\cdot 365d/y}{EF_x\left[\left(\frac{IFS_{adj}\cdot CSF_o}{10^6mg/kg}\right) + \left(\frac{SFS_{adj}\cdot ABS\cdot CSF_o}{10^6mg/kg}\right) + \left(\frac{InhF_{adj}\cdot CSF_i}{VF^a}\right)\right]}$$

Equation 3-2: Direct Exposures to Noncarcinogenic Contaminants in Residential Soil

$$C(mg/kg) = \frac{THQ \cdot BW_c \cdot ED_x \cdot 365 d/y}{EF_x \cdot ED_c \left[\left(\frac{1}{RfD_o} \cdot \frac{IRS_c}{10^6 mg/kg} \right) + \left(\frac{1}{RfD_o} \cdot \frac{SA_c \cdot SL \cdot ABS}{10^6 mg/kg} \right) + \left(\frac{1}{RfD_d} \cdot \frac{IRA_c}{VF^a} \right) \right]}$$

Equation 3-3: Direct Exposures to Carcinogenic Contaminants in Industrial Soil

$$C(mg/kg) = \frac{TR \cdot BW_a \cdot AT365d/y}{EF_o \cdot ED_o \left[\left(\frac{IRS_o \cdot CSF_o}{10^6 mg/kg} \right) + \left(\frac{SA_a \cdot SL \cdot ABS}{10^6 mg/kg} \right) + \left(\frac{IRA_a \cdot CSF_i}{VF^a} \right) \right]}$$

Equation 3-4: Direct Exposures to Noncarcinogenic Contaminants in Industrial Soil

$$C(mg/kg) = \frac{THO \cdot BW_a \cdot ED_o \cdot 365 d/y}{EF_o \cdot ED_o \left[\left(\frac{1}{RfD_o} \cdot \frac{IRS_o}{10^6 mg/kg} \right) + \left(\frac{1}{RfD_o} \cdot \frac{SA_a \cdot SL \cdot ABS}{10^6 mg/kg} \right) + \left(\frac{1}{RfD_d} \cdot \frac{IRA_a}{VF^a} \right) \right]}$$

Footnote:

^{*}Use VF for volatile chemicals (defined as having a Henry's Law Constant [atm-m³/mol] greater than 10⁻⁶ and a molecular weight less than 200 grams/mol) or PEF for non-volatile chemicals.

Tap Water Equations:

Equation 3-5: Ingestion and Inhalation Exposures to Carcinogenic Contaminants in Tap Water

$$C(ug/L) = \frac{TR\cdot AT\cdot 365d/y\cdot 1000ug/mg}{EF_x[(IFW_{adj}\cdot CSF_o) + (VF_y\cdot InhF_{adj}\cdot CSF_i)]}$$

Equation 3-6: Ingestion and Inhalation Exposures to Noncarcinogenic Contaminants in Tap Water

$$C(ug/L) = \frac{THQ \cdot BW_a \cdot EDr \cdot 365 d/y \cdot 1000 ug/mg}{EF_x \cdot ED_x \left[\left(\frac{IRW_a}{RfD_o} \right) + \left(\frac{VF_y \cdot IRA_a}{RfD_i} \right) \right]}$$

Air Equations:

Equation 3-7: Inhalation Exposures to Carcinogenic Contaminants in Air

$$C(ug/m^{2}) = \frac{TR \cdot AT365 d/y \cdot 1000 ug/mg}{EF_{z} \cdot InhF_{adj} \cdot CSF_{i}}$$

Equation 3-8: Inhalation Exposures to Carcinogenic Contaminants in Air

$$C(ug/m^3) = \frac{THQ \cdot RfD_i \cdot BW_a \cdot ED_x \cdot 365d/y \cdot 1000ug/mg}{EF_x \cdot ED_x \cdot IRA_a}$$

SOIL-TO-AIR VOLATILIZATION FACTOR (VF)

Equation 3-9: Derivation of the Volatilization Factor

$$VF(m^3/kg) = \frac{(LS \cdot V \cdot DH)}{A} \cdot \frac{(3 \cdot 14 \cdot \alpha \cdot T)^{1/2}}{(2 \cdot D_{el} \cdot P_e \cdot K_{es} 10^{-3} kg/g)}$$

where:

$$\alpha = \frac{D_{ei} \cdot P_a}{P_a + [(\rho_s) (1 - P_s) / K_{as}]}$$

Param	peter Definition (units)	Default
VF	Volatilization factor (m²/kg)	
LS	Length of side of contaminated contaminated area (m)	45
V	Windspeed in mixing zone (m/s)	2.25
DH	Diffusion height (m)	2
A	Area of contamination (cm²)	20,250,000
De	Effective diffusivity (cm²/s)	$D_i(P_a^{a.so}/P_i^a)$
P.	Air filled soil porosity (unitless)	P ₁ - 0 6
Pt	Total soil porosity (unitiess)	1-(B/p _a)
0	Soil moisture content (cm²-water/g-soil)	0.1
В	Soil bulk density (g/cm²)	1.5
Pa	True soil density or particle density (g/cm²)	2.65
K_	Soli-air partition coefficient (g-soll/cm²-air)	(H/K _a) x 41 (41 is a conversion factor)
T	Exposure interval (s)	7.9 x 10°
D,	Diffusivity in air (cm²/s)	Chemical-specific
Н	Henry's Law constant (atm-m²/mol)	Chamical-specific
K	Soil-water partition coefficient(cm²/g)	K _{ex} x OC
k.	Organic carbon partition coefficient (cm²/g)	Chemical-specific
oc	Organic carbon content of soil (fraction)	0.02

SOIL SATURATION CONCENTRATION (C_)

Equation 3-10: Derivation of the Soil Saturation Limit

$$C_{aat} = \frac{S}{\beta} \left(K_d \beta + P_w + H' P_a \right)$$

P	arameter	Definition (units)	Default
C,		saturation concentration (mg/kg)	-
K,	Soil-v	water partition coefficient (mg/kg-soil per mg/L-water)	Chemical-specific, or K _{sc} x OC
K,	e Organ	nic carbon partition coefficient (mg/kg-OC per mg/L-water)	Chemical-specific
0	C Organ	nic carbon content of soil (kg-OC/kg-soil)	Site-specific or 0.02
s	Solut	olity in water (mg/L-water)	Chemical-specific
В	Soil d	iry bulk density (kg-soll/L-soll)	Site-specific or 1.5
P.	Water	r filled soil porosity (unitiess)	P _t · P _a
H	Henn	/'s Law constant (unitless)	H x 41, where 41 is a units conversion factor
н	Henry	r's Law constant (atm-m²/mol)	Chemical-specific
P.	Air-fill	ed soil porosity (unitless)	P ₁ - 0 B
0		ige unsaturated zone soil volumetric content (L-water/L-soil)	O_B/t_
0_	gravin	ige unsaturated zone soil netric water content ster/kg-soil)	Site-specific or 0.1
r,	Densi	ty of water (kg-water/L-water)	1 kg/L
Pı	Total	soil porosity (unitiess)	1 - B/p.
ρ.	True	soil density or particle density (kg-soil/L-soil)	2.65 kg/L

ATTACHMENT A PHYSICAL-CHEMICAL CONSTANTS USED IN CALCULATING SOIL PRGs FOR VOLATILE ORGANIC CHEMICALS

	MW (g/mol)	Henry's Law (atm-m^3/mol)	Diffusivity in Air (cm^2/s)	Kec (ml/g)	Water Solubility (mg/l)	References
Acetone	58	0.000021	0.100	2.2	1000000	
Acrytonitrile	53	0.000088	0.110	0.9	79000	
Ammonia	17	0.000320	0.260	3.1 65.0	530000 1800	
Benzene .	78 130	0.005500 0.000051	0.087	50.0	3300	
Benzyl chloride	140	0.000290	0.070	14.0	10000	
Bis(2-chloroethyl)ether	170	0.000110	0.063	61.0	1700	
Bis(2-chloroisopropyl)ether	120	0.000200	0.089	1.2	22000	1,2
Bis(chloromethyl)ether	160	0.001600	0.080	100.0	4700	2,4
Bromodichloromethane Bromoethene (Surrogate = Bromomethane)	108	0.006200	0.100	130.0	18000	2,4
Bromemathane	95	0.006200	0.100	130.0	18000	
1,3-Butadiene	. 54	0,180000	0.098	120.0	740 2900	
Carbon disuffide	76	0.012000	0.110 0.080	54.0 110.0	760	
Carbon tetrachloride	150	0.024000	0.060	110.0	,	-,0
Chlorine dioxide						
Chloroacetaldehyde	150	0.003500	0.072	160.0	470	2,3
2-Chloroacetophenone (Surrogate = Chlorobenzene)	110	0.003500	0.072	160.0	470	2,3
Chlorobenzene C-Chloro-1,3-butadiene	88	0.032000	0.110	50.0	860	
-Chlorobutane (Surrogate = 2-Chloro-1,3-butadiene)	93	0.032000	0.110	50.0	680	
Chlorodifluoromethane (Surrogate = Dichlorodifluoromethane)	120	0.100000	0.080	58.0	280	1,2,4
I-Chloroethyl vinyl ether					8000	404
Chloroform	120	0.003800	0.089	31.0	8200 8200	1,2,4
Chloromethane	51	0.024000	0.110 0.080	35.0 51.0	2700	1,2,4 1,2
2-Chloropropane	79 127	0.002300 0.003500	0.072	180.0	470	2,3
o-Chlorotoluene	70	0,240000	0.091	840.0	20	1,2
Crotonaidehyde (Surrogate = Methyl methacrylate)	120	0.006400	0.075	220.0	150	2,3
Curnene (Surrogate = Ethylbenzene) I 2-Dibromoethane	190	0.000320	0.073	28.0	3400	2,3
t_2-Dichlorobenzene	150	0.001900	0.130	1100.0	100	2,3
t.3-Dichlorobenzene	150	0.001900	0.130	1200.0	120	2,3
,4-Dichlorobenzene	150	0.001600	0.130	1200.0	79	2,3
1,4-Dichloro-2-butene (Surrogate = 2-Chloro-1,3-butadiene)	122	0.032000	0.110	50.0 58.0	660 . 280	1,2 1,2,4
Dichlorodifiuoromethane	120	0.100000 0.004300	0.080 0.091	30.0	5500	2,3
1,1-Dichloroethane	89	0.001200	0.091	14.0	8700	2,3
t.2-Dichloroethane (EDC)	97	0.150000	0.079	65.0	400	2,3
I,1-Dichloroethylene I,2-Dichloroethylene (trans)	97	0.006600	0.079	59.0	6300	. 2,3
1,2-Dichloroethylene (mixture)	97	0.006600	0.079	59.0	6300	2,3
1,2-Dichloropropane	110	0.003600	0.080	51.0	2700	1,2,4
I_3-Dichloropropane	110	0.001300	0.080	48.0	2800	1,2
1,3-Dichloropropene	110	0.001300	0.081	48.0	2800	1,2
Dicyclopentadiene			0.400		1000000	1,2
Dimethylamine	45 88	0.000090 0.000011	0.120 0.085	2.2 3.5	430000	1,2
1,4-Dioxane	93	0.000011	0.088	3.5	80000	1,2
Epichlorohydrin Ethyl acrylate (Surrogate = Methyl methacrylate)	100	0.240000	0.091	840.0	20	1,2
Ethylbenzene	110	0.007900	0.075	220.0	680	2,3
Ethylene oxide	44	0.000076	0.130	2.2	1000000	1,2
Ethyl chloride	65	0.011000	0.100	15.0	5700	2,3
Ethyl ether ·	74	0.000013	0.070	14.0	10000	1,2,4
Ethyl methacrylate (Surrogate = Methyl methacrylate)	120	0.240000	0.091	840.0	20	1,2
lydrogen sulfide		0.000000	0.440	0.9	79000	1,2
Methacrylonitrile (Surrogate = Acrylonitrile)	93 74	0.000088 0.000021	0.110 0.100	2.2	1000000	1.2
Methyl acetate (Surrogate = Acetone)	100	0.240000	0.091	840.0	20	1.2
Hethyl acrylate (Surrogate = Methyl methacrylate)	85	0.002600	0.100	8.8	13200	2.3
Methyl ethyl ketone	72	0.000027	0.090	4.5	270000	2,3
Methyl styrene (mixture) (Surrogate = Styrene)	119	0.002300	0.071	360.0	300	2,3
Methyl styrene (alpha) (Surrogate = Styrene)	119	0.002300	0.071	360.0	300	2,3
Vitrogen dioxide						
!-Nitropropane						
Polynuclear aromatic hydrocarbons				****		
Acenaphthene	150	0.001200 0.000034	0.064 0.058	4600.0 13009.0	4 0	2,3 2,3 2,3
Anthracene Fiuorene	180 170	0.000084	0.081	7900.0	2	2,3
Nachthalene	130	0.001300	0.069	1300.0	31	2,3
Phenanthrene	180	0.000040	0.058	14000.0	1	2,3
Propylene oxide	58	0.000010		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
Styrene	100	0.002300	0.071	360.0	300	2,3
,1,1,2-Tetrachioroethane	170	0.000380	0.073	54.0	2900	1,2
1,1,2,2-Tetrachloroethane	170	0.000500	0.073	220.0	2900	2,3
Tetrachloroethylene (PCE)	170	0.023000	0.072	0,00	150	2,3
Tetrahydrofuran	72	0.000110	0.089			2,3
Coluene	92	0.008600	0.078	260.0	520	2,3 2,3 2,3 1,2
1,2,4-Trichlorobenzene	180 130	0.002300 0.002800	0.062 0.080	9200.0 150.0	30 950	1,2
t,1,1-Trichloroethane t,1,2-Trichloroethane	130	0.001200	0.080	56.0	4500	2,3 2,3
Frichlargethylene (TCE)	130	0.008920	0.081	130.0	1000	2,3
Frichlorofluoromethane	137	0.097000	0.087	160.0	1100	1,2,4
1.1,2-Thichloropropane (Surrogate = 1,2-Dichloropropane)	147	0.003600	0.080	51.0	2700	1,2
1.2.3-Trichloropropane (Surrogate = 1,2-Dichloropropane)	147	0.003600	0.080	51.0	2700	1,2
	146	0.001300	0.081	48.0	2800	1,2
1,2,3-Trichloropropene (Surrogate = 1,3-Dichloropropene)				400.0	1100	1,2,4
	186	0.058000	0.087	160.0	1100	
1,1,2-Trichloro-1,2,2-trifluoroethane (S = Trichlorofluoromethane) Triethylamine (Surrogate = Dimethylamine)	186 86	0.000090	0.120	2.2	1000000	1,2
1,1,2-Trichloro-1,2,2-trifluoroethane (S = Trichlorofluoromethane) Triethylamine (Surrogate = Dimethylamine) Vinyl chloride	186 86 63	0.000090 0.700000	0.120 0.110	2.2 57.0	1000000 1100	1,2
1,1,2-Trichloro-1,2,2-trifluoroethane (S = Trichlorofluoromethane) Triethylamine (Surrogate = Dimethylamine) Vinyl chloride m-Xylene	186 86 63 110	0.000090 0.700000 0.006900	0.120 0.110 0.087	2.2 57.0 240.0	1000000 1100 200	1,2 2,3 2,3
1.2.3-Trichloropropene (Surrogate = 1.3-Dichloropropene) 1.1,2-Trichloro-1.2,2-trifluoroethane (S = Trichlorofluoromethane) Triethylamine (Surrogate = Dimethylamine) Vinyl chloride m-Xylene o-Xylene p-Xylene	186 86 63	0.000090 0.700000	0.120 0.110	2.2 57.0	1000000 1100	1,2

20
080
mucker
5.1.5
Soals:
lation (
Semed
Ilnary F
Prefit
S wold
A Re
-

(PRGs)	25555555555555555555555555555555555555	252955555 252955555		**************************************
S 0	7.80 2.20 2.20 2.20 3.20 3.20 3.20 3.20 3.2	2 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	2001 100	3.8E-00 1.5E+01 1.5E+01 1.8E-02 1.8E-03 1.5E+03 1.1E+03 1.1E+03 1.1E+03
	7. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	2002222500 595825558	25	4.7E+01 inc 8.2E+02 inc 8.2E+03 inc 8.2E+0
REMEDIAL Ambert	9.4 7.3 1.0 1.0 1.1 1.1 1.1 1.1 1.1	3.16.40 3.16.40 3.76.40 3.76.40 3.76.40 3.76.40 3.76.40	1.1E+00 1.0E+00 2.3E+01 7.3E+01 1.0E+02 1.0E+02	4.76-01 2.76-01 2.76-03 3.36-02 3.36-02 3.16-02 3.16-02 3.26-03 1.56-03 1.56-03 1.56-03 1.56-03 1.66-0
F X La		25255222		
SES EST MENTARY ELIMINARY SOIL (mpling	2.25 + 0.2 1.85 + 0.2 1.16 + 0.4 1.16 + 0.4 1.85 + 0.4 1.55 + 0.3 1.55 + 0.3	2.4E+05 1.0E+05 6.0E+05 6.0E+02 1.0E+02 1.0E+02 1.0E+02 1.0E+02 1.0E+02	3.4E+43 3.4E+43 3.4E+43 3.4E+43 3.4E+43 4.4E+43 4.4E+43 4.4E+43 4.4E+43 4.4E+43 4.4E+43 4.4E+43 6.6E+63 6.6	2.05.03 2.05.03 2.05.03 2.05.03 2.155.03
	25555555	2 2 2 3 5 2 2 2 3 5	222222222222222	22222525 2223252222
URPC SOIL SATURATION PR Residentia Soil (marka)	2.00 + 0.	2.2E-01 2.2E-0	33E-03 31E-04 31E-04 31E-04 4.8E-02 1.8E-02 1.8E-04 3.1E-04	22 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
PLANNING P OLATION EMECANCER PRIG INC. HONCANCER PRIG SER RS CONTAMINANT	1988888888		24444444444444444444444444444444444444	Artimory penloxide Artimory todasslum lartrate Artimory tetroxide Aramile Aram
FACTOR	00000000000000000000000000000000000000	0 7.8E+03	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
SOIL F.				
MITHDRAWN JES (mg/kg-d)	2.00 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2.0E-04 1.0E-04 1.0E-05 1.0E-0	╒┥┋┋┋	250
TY VALU	8.7E-03	4.8E+00 I 2.4E-01 I 6.0E-02 I	5.7E-03 [2.5E-021 1.5E-011 2.2E-011
TOXIC)	2.0E-03 1.0E-03 1.0E-0	2.2.2.1.1.1.0.0.0.0.0.0.0.0.0.0.0.0.0.0.	2.2.7.7.00.00	24
Key: FIRS	8.7E-03.1	6.4E-01 i 8.1E-02 i 1.7E+01 j	5.75-03	2.5E-021 1.8E+001 2.2E-01 h

_
3
\$
ğ
8
=
¥
ş
ā
-
8
100
8
5
흄
묶
ē
5
2
Ē
Ē
Ŧ
2
-
5
₫
8
2
2
101

MIT 'no" < 100X ca **= io < 10X ca T REMEDIAL GOALS (PRGs) Ambient Ar Tsp Weier (upin/3) (upin/3)	1 E+03 n 1 E+04 n 1 E+04 n 1 E+05 n n n n n n n n n	2.1E+01 nc 3.3E+03 1.1E-02 ca* 1.0E-01 8.1E-02 ca* 1.0E-01 6.2E+00 nc 8.5E+00 7.3E+01 nc 1.8E+02 7.3E+01 nc 1.8E+02 7.3E+02 nc 1.8E+02 8.9E+03 nc 3.7E+03 0x 7.3E+02 nc 3.7E+03 0x 7.3E+03 nc 7.3E+03 0x 7.3E+03 nc 3.7E+03 0x 3.7E+03 nc 3.7E+03 0x 3.7E+03 nc 3.7E+03 0x 3.7E+03 nc 3.7E+03	5 05 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
URPOSES SOIL SATURATION MENTICELLING LIN Residentia Residentia Soil (marks) Soil (marks) Soil (marks)	200-604 no control of the control of	0.00	3.3E+17 Inc. 3.3E+17 Inc. 3.3E+17 Inc. 3.3E+17 Inc. 1.0E+17 Inc. 2.2E+17 Inc. 2.3E+17 Inc. 2.3E+17 Inc. 2.3E+17 Inc. 3.4E+17 Inc. 3.4E+
PLANNING PUI	Benefin Benefin Benzaldehyde Benzaldehyde Benzaldrive Benzaldrive Benzaldrive Benzaldrive Benzola acid Benzol	mide) (fher	(PEA 1994)
FOR EXTRAPO SOIL FACTOR O SIGN OF CARS (m*3/kg)	0000-00000000	22 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
ICITY VALUES ISF IRD IMMENDED ISF IRD	2.3E-021 1.7E-051 1.7E-051 1.2E-000 1.2E-000 1.4E-021 1.4E-027	22 6.28 02 2.06 02 0.28 02 0	
Key: EIRS h=HEAST TOXIC SF ORD Managed (managed)	2.9E-02 2.3E+02 1.3E+02 1.3E+02 1.3E+00 1.7E-01 1.1E+00 1.1E+0	8.2E-02 2.0E-02 1.1E-01 2.0E-02 1.2E-03 1.2E-03 1.2E-03 1.2E-03 1.2E-03 1.2E-03 1.0E-03 1.0E-0	3.5E-03 h 1.3E-01 2.0E-02 h 1.3E-01 1.3E-01 h 1.0E-01 1.3E-01 h 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01 1.0E-01

	(PRGs)	1.7E-01 cs 5.2E-02 cs 7.3E+02 nc 3.7E+03 nc	2.2E-02 no	8.5E+04 RE 1.5E+09 CS 1.5E+09 CS	1.25+02 no 7.35+03 no 3.75+02 no 1.85+03 no 2.95+01 no 1.85+02 no 1.85-02 no	1.4E-03 nc 1.9E-03 nc 1.9E-03 nc 1.9E-03 nc 1.5E-03 nc 1.5E-03 nc 1.5E-03 nc 1.5E-03 nc 1.5E-03 nc 1.5E-03 nc 1.5E-03 nc 1.5E-03 nc 1.5E-03 nc
C < 10X CB	ALS	3.7	25.5	8.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1	22.1.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.	200 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
1	Ø 36	882 2	2222222222	2888888888888		5 525
nc= < 100X cs	Amberi (usim	1.76.02 5.26.03 7.36+01 2.16.01	7.36-407 3.16-07 1.86-401 2.56-402 7.36-402 7.36-403 7.36-403 7.36-403 7.36-403	5.16.404 1.16.402 1.26.402 2.96.402 2.76.01 3.76.01 1.06.402 6.16.01	7.3E+00 7.3E+00 7.1E+00 7.1E+0	3.1E-03 9.4E-00 6.0E-03
MIT	> L ₉	8888	222252225	3535555555		5952 4555455
SES	ELIMINAR Industria Sal (mg/k	4.7E+00 1.5E+00 1.4E+04 1.7E+05	1.4E+03 3.8E-01 2.7E+03 7.1E+02 1.0E+05 1.4E+04 1.4E+04 6.1E+02	2.5.4.02 2.3.3.4.00 2.3.3.4.00 2.4.1.6.	1.0E+05-05-05-05-05-05-05-05-05-05-05-05-05-0	8.38E-02 2.3E-02 2.3E-02 1.0E-03 6.5E-03 1.0E-04 3.4E-04 1.4E-04
	A L	2255		# 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	2 2 2 2 2 2	2525 2222222
URP	Residentia Soil (mg/kg	1.1E+00 3.4E-01 1.3E+03 7.7E+03		3.5E+02 7.7E+03 7.7E+03 7.7E+03 8.7E-01 1.8E+01 2.5E+03 3.3E+02 4.7E+02 4.0E+01	1,45-40 1,35-40 2,05-40 3,35-40 5,25-40 3,86-40 2,06-40 2,06-40	2.8E+02 1.6E-02 1.3E+02 1.3E+03 3.8E+03 2.8E+03 5.9E+03 5.9E+03 1.3E+03
PLA	CONTAMINANT	Chortanii Chortanii Chortanii Chorta	Chlorocetic acid Chlorocetic acid Chlorocetic acid Chloroberzale	200442997700 2550044299700055		Copper cyanide
OR	ACTO!		2.9E+03 2.9E+04 2.9E+03 2.9E+03	1.8E+03 2.8E+03 2.8E+03 1.4E+04		1.08 1.08 1.08 1.08 1.08 1.08
H B	PA PARS	000000				
E	0 > 0 U	0000-	0-0-000			000000000000000000000000000000000000000
MITHDRAWN	MES (mg/kg-0)	2.0E-05 5.7E-05	48688888888888888888888888888888888888		2.	7. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.
e:ECAO x=	ITY VAL	1.3E+00	2.76-01	8.1E-02 5.8E-01 7.5E-02 1.8E-02 1.1E-02		x 1.9E+00 h 8.4E-01
S h=HEAST	TOXIC ORID (Mg/kg-d)				2.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
Key: FIR	OSF 11(mg/kg-d	4.0E-01 1.3E+00	2.7E-01	6.1E-03 5.8E-01 5.8E-01 1.8E-02 1.8E-02 1.1E-02		1.9E+001

8/01/94]
5
Smucke
8
Goels:
Remediation
Preliminary
×
Region
EPA

REMEDIAL GOALS (PRGS) Ambent Ar Tap Water (ug/m ³) (ug/m	1.8E+02 Inc. 1.8E+03 Inc. 1.8E+04 Inc. 1.8E+
SES MARKEN FEW SOIL (MINARY REW SOIL (MUSICE)	- 1
	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
PLANNING PURP TON CHANGER PRIG TO-NONCANCER PRIG SEET SOIL SATURAT CONTAMINANT Residentia Soil (mg/k)	Hydrogen cyanide Polassum sher cyanide Polassum cyanide Sociam sher cyanide Sociam cyanide Sociam cyanide Sociam cyanide Sociam cyanide Sociam cyanide Cyclotexanone Cyclotexanone Cyclotexanone Cyclotexanone Cyclotexanone Cyclotexanone Cyclotexanone Cyclotexanone Darina Darin
FOR MINDRAWA THROUTE EXTRAPOLA JES SOIL FACTORS (MRTD O SIGN OF MARKE) CABS (m*3/kg)	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
Key: EIRS h=HEAST e=ECAO x= TOXIC)TY VAL: oSF oRID ISF	2.4E.01 5.0E.02 4.0E.02 5.0E.02 5.0E.02 5.0E.02 5.0E.02 5.0E.02 5.0E.02 5.0E.02 5.0E.02 5.0E.03 5.0E.04 7.0E.03 7.0E.04 7.0E.03 7.0E.04 7.0E.0

1	_
3	Ē
1	2
Š	5
2	ę
-	=
•	9
1	ğ
i	Ē
Ġ	5
-	i
	Ö
:	
ì	Ĭ
ŀ	2
7	Ē
4	5
3	문
7	B
Į	Ę
è	č
1	-
1	ē
1	Ē
ŧ	Ē
å	F
5	ē
•	3
i	5
3	ì
8	
3	C
0	i

	PRGs	Tap Water (ug/l)	205	102	2.9E+04 nc	E-05 CB	.3E+02 Inc	+02 IC	3E+00 nc	SE-02 no		0E-02 CB	35-03		11-03 G	E+02 nc	2E+01 nc	5 5 5	+03 Inc	5 5 5	E+01 nc	200	8.9E-02 ca	36+01	7-10-12	E+02 nc	E+00	E+02 mc	E-02 cs	100 E	E-03 ca	E-03	102	E+04	100 TC	E+02	L' W. I. W. L.
enc < 10X cs	GOALS (PRGs	Ш	2.1	4.0	2.9		7.3	7.3	6.	3.5			1	2		7.3	2.2	37	3.7	3.7	.5	5:5	1.6	7.3	7	7.3	21	9 2	70	36	8.	7.	3.7	2		2.7	-
nc= < 100X cs "==;		Ambient Ar (up/m/3)	2.1E+01 mg	4.0E+01 mg	2.8F+03 2.8F+03 75	2.9E+02 no	7.3E+01 no	7.3E+01 no	7.3E-01 no	2.1E-02 Inc	7.3E+00 nc	9.06-43	7.3E-04 Ca	1.95-03	3 15 0	7.3E+01 nc	2.2E+00 IIX	3.75.405	3.7E+02 nc	3.75-017	1.5E+00 m	1.5E+00 F	9.9E-03	7.3E+00 IIX	3.75+00 0	7,3E+01 IN	6.1E-01 Ca	9.1E+01 Inc	_	7 AE-04 IC		7.2E-04 CS	3.76-0	7.3E+00 IIV	1.8E-01 IN	7.3E+01 IR	11 11 11 11
MIT	ELIMINARY REMEDIA		Tight make	2		8 2	2	5 5		3 2	2	8 8	8	8	8 2	2	5 5			2 2	2	2 2	2 8	2	3 2	2	88	100	8	28	5	8 2	2 2	2	5 5	28	2
SES	IMINAR	Soll (mg/kg)	3.9E+03	7.5E+03	1.0E+03	5.5E+04	1.46+04	1.4E+04	1.4E+02	3.2E-01	1.4E+03	3.35.+06	2.1E-01	7.3E-01	8.8E+04	46.45	4.15+02	1.0E+05	6.8E+04	8.85+03	2.7E+02	2.75.402	2.8E+00	1.4E+03	6.8E+02	1.45+04	4.8E+01	1.7E+01	2.4E+00	2.2E-01	2.4E-01	2.1E-01	6.05+03	1.4E+03	3.4E+01	1.4E+04	
	PREL	ofke)	02 nc	02 nc	04 Inc	03 cs 03 nc	93 nc	03 nc	200	02 Inc	-05 Inc	50	02 Cs	01 03	93 03	-03 Inc	2 2 2	05 max	03 Inc	2 2 2 2	0 10 10	20 20	-01 Ca	-02 Inc	200	-03 mc	E+01 CB	-03 mc	01 Cs	02 CS	-02 CG	92 68	02 Inc	102 Inc	-00 mc	-03 nc	
Solt satur		Residential Soil (mg/kg)	3.7E+02	7.25	3.7E+0.	8.5E-05 5.2E+03	1.35+	1.3E+03	1.35	8.35	13	7.16	4.8E	1.7E-01	6,5E+03	1.35	3.95+0	1.06-	6.5E-	1.3E+02 6.5E+00	2.6E+0	2.8E+0	6.5E-0	1.3E+0.	6.5E+01	1.35	1.87	1.6E-	5.6E-0	525	5.5E-02	4.8E-02	6.5E+02	1.3E	3.3E	1.3E	
P P P R SERVER																																					
PLANNING PURPC	INANT		other					P				92																									
N N N N N N N N N N N N N N N N N N N	CONTAMINAN		e glycol, monobutyl ether e glycol, monoethyl ether	- Constant	Thelate	(edu	enzigon enem malhumboenhoode	I de la company	hmvhenzirina		ine	ne hydrochlo	zidine	azine	nemide	2	entry phenol	te te	halate Aband aban	Ne Ne	94	2	xtrue			erte			Bzine								
LA	3		hyene glyco	Moramide	M Phihalate	nzogual (Avenge	penzinon	ethor	Silvoste Dimethoriz	ethylamine	Omelhylanline		Omethylben		Dimethyllon	Dimethylphe		ethyl phthala	of the teneohit	Uninteenze	Dinitrobenze	Omtropheno	trotoluene m		2860	Octyl phithelate	Moname	henylamine	Dipherythyd	rect black 38	ct blue 6	ect prown 35 suffoton	-Dithiane	5	dosullan	othail	
OR P	ACTORS	(m*3/kg)	000			58		3E	260	.2E+04 Dim	2	24	33	- 6	N.W	2.4	37.6	E.	E .	1.3	**	24	6	2,4		40 100	Ţ	8	1.2	35	O		1.4		5		
F		ABS	0.10			_	_	_		0.10	_		_		_			_	_	_					_	0.10	000	_	_	_	-	_	_		_		
HDRAWN	S.	(mg/kg-d)	5.7E-03 × 0	J.		6.0E-02 r		20E-02	200	6.7E-08 x	1	0				2.0E-02 0	1.00-03				Π.	205-301			1.0E-03 r 0	I	1.0E-02 r 0	2.5E-02 r 0	2 7 F. A. S. L. O		OK 	.0E-05 0			IJ	3.0E-04 F 0	1
ECAO x=WITH	VALUES	13F 11(mg/kg-d) (m		4 9E IN	Ц.	╜			1.4E-02	Н	+	8E-01	2E+00 r	3.7E+01 ×	Н								6.8E-01 r			4 E 705	1		(,/5-01 /	Н	E+00	+					
HEAST 61EC		oRiD (mg/kg-d) 1/(m	5.7E-03 h		H	廿	F-02	2.0E-02	+	5.7E-08 r	#	5	6)	Ť	힏	0E-02	1.0E-03	E+01 h	100	E-04	24 P	\pm	H	7E-03 IA	1.0E-03	_	Ł	2.5E-02	2 25-63		6		E-02	E-03	E-05 h	3.0E-04	
:: 			5.0	1 2F-M3 6 0	1		7.7	2	1.4E-02 h	Н	+	01 h	400+	3.7E+01 ×	#	2,5	1,0	6		F	***	+		107	2	‡	1	H	+	님	U 00+	+	1:0	7.0	5.0	3.0	
Key:		0SF 1/(mg/kg-d		-	Ш				\perp	1 1													6.8E-0			2 4		200	9	8.8E+00	0.15	9.3					

GOALS (PRGs)	2.7.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2
IN LIEE	10 10 10 10 10 10 10 10
SES MARY REMEDIA Includia Sol (marks) (9)	3.3.2.2.4.0.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2
	1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
PLANNING PURP CONTAMINANT CONTAMINANT Residential Soil (market)	Epichkrofyddin 1.2-Epoxyblane EPTC (S-Ehri dipropythiocarbamele) Ehrebron (2-chloroethyd prosphonic add) Ehrebron (2-chloroethyd prosphonic add) Ehriboxyethanol acetale Ehriboxyethanol acetale Ehry ac
ALUES SOIL FACTORS FIND O SIGN OF MANAGED IN THE STRAPOLATION OF SIGN	4.2E-03 2.9E-04 1 0.10 3.0E+04 2.5E-03 0.10 0.10 0.10 3.0E+04 1 0.10 3.0E+04 1 0.10 3.0E+04 1 0.10 3.0E+04 1 0.10 0.10 0.10 0.10 0.10 0.10 0.10
Key: IsIR 5 heHEAST evECAO xer TOXIC TY VAL. OSF ORID ISF	4.8E-03 2.0E-03 4.3E-03 2.0E-03 4.3E-03 2.0E-03 4.3E-02 6.0E-04 4.3E-02 6.0E-04 6.0E-0

	LS (PRGs)	23.7.7.6.00	3.7E+01 III 9.1E+02 III 1.8E+02 III 3.7E+01 III
	GOA S⊄	######################################	2.7E+00 1.E+01 in 1.E+01 i
SES	ELIMINARY R Industrial Soil (mg//gi)	2. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	6.8E+02 nc 1.7E+04 nc 3.4E+03 nc 6.8E+02 nc
URPC	Residentis Soil (mg/k	2.516-62 2.516-62 2.516-63 2.516-	1.6E+01 nc 1.6E+01 nc 3.3E+02 nc 6.5E+01 nc
PLANNING P		HCH (eiche) HCH (cethe) HCH (demme) Indene HCH (gemme) Indene HCH (gem	Weihdaihon Weihomy Weihowychor 2-Meihowyeihanol
FOR THOMAS	S SOL FACTOR V V (mp/mg-d) CABS (m-3/mg)	2.00	1.0E-03 (r 0 0.10 2.5E-02 (r 0 0.10 5.0E-03 (r 0 0.10 5.7E-03 (l 0 0.10
E R S E E C A S E E C A S E E C A S E E C A S E E C A S E E C A S E E E E E E E E E E E E E E E E E E	TOXICITY VAL	00 1 3 0 0 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1.0E-03 2.5E-02 5.0E-03 1.0E-03 h
Key:		1.3E+00 1.3E+00 1.4E-02 1.4E-02 1.1E-07 1.1E-07 1.1E-07 1.1E-07 1.1E-07 1.1E-07	

		2 2 2 2 2		8 2 8 8 8 8 8	595595555		20 20 20 20 20 20 20 20 20 20 20 20 20 2
	GOALS (PRGs	(ur 7.3E+0 1.5E+0 6.1E+0 1.8E+0	3.7E-01 3.7E-01 3.7E-01 3.7E-02 3.7E-02	2.E-01 2.E-01 5.2E-01 3.7E+00 4.3E+00	2.9E+03 2.9E+03 2.9E+03 2.0E+00 2.0E+00 1.8E+03 1.8E+03	6.0E-01 4.3E-02 1.8E-02 8.1E-02 3.7E-02 1.3E-02 1.3E-02 1.3E-03 1.7E-03 1.7E-03 1.7E-03	3.15+02 7.3(5+02 5.5(5+04) 5.6(5+04) 3.7(5+03 2.7(5+03)
		S 2 2 2 2	8828888	2288828	523225222	22222222222	2 22 2
	REMEDIAL Ambient	1.5E-01 3.7E-03 1.6E-03		2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.	2.05-02.02 1.05-03.03 0.35-03 10 2.05-03 10 2.05-03 10 3.15-03 10 1.85-02 10 1.85-02 10 1.85-02 10	4.2E+01 2.6E+02 3.1E+03 9.1E+01 7.3E+00 1.3E+00 1.3E+00 1.3E+00	3. fe-02 4.0E-03 5.5E+00
	- 12	8 5255	5522222	288828	282282222	2222222222222	
SES	ELIMINARY Industral	3.8E+04 2.1E+02	8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	3.9E+00 7.6E+00 1.5E+01 4.1E+01 6.8E+03	2. 1. 1. 2. 2. 1. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	3.3.8.6.4.1.1.1.3.8.6.6.4.3.3.8.6.4.1.3.8.6.4.1.3.6.4.	3.9E-04 (4.1)
	Z C	2822	5522222	252225	282282222	222222222222	
		1.3E+02 1.0E+02 1.0E+04	2.55 2.55 2.55 2.55 2.55 2.55 2.55 2.55	3.7E-01 1.8E-00 3.4E+00 6.7E+00 6.5E+02 4.9E+00	4.2E+03 4.0E+03 5.2E+03 1.3E+01 1.3E+01 3.3E+03 3.3E+03 3.3E+03 3.3E+03	1.3 E+02 3.9 E+02 3.9 E+03 1.0 E+03 3.9 E+03 3.9 E+03 3.9 E+03 5.5 E+03 5.5 E+03 5.5 E+03 5.5 E+03 7.5	8
	RS CONTAMINANT Residential Res	2 Methoxyethan 2-Methoxy-5-rit Methy acetate Methy acryate	2-Methyleniline to-cumpine) 2-Methyleniline hydrochloride 2-Methyl d-chlorophenoxyacetic acid 4-(2-Methyl-4-chlorophenoxyacetic acid (MCPB) 2-(2-Methyl-4-chlorophenoxy) bulync acid (MCPB) 2-(2-Methyl-4-chlorophenoxy) propionic acid 2-(2-Methyl-1-4-chlorophenoxy) propionic acid Methylenic acid (MCPP)	Methy A.	Methylethylethylethylethylethylethylethyl		Nickel and compounds "CAL-Modified PRG" (PEA, 1994) Nickel refinery dust Nickel refinery dust Nickel subsuffide Nifratie
F	IL FACTO	00000					
	JES SO	2.0E-03 r 1.0E+00 r 3.0E-02 r	2.000 00 00 00 00 00 00 00 00 00 00 00 00	7.0E.04 r 1.0E.02 r 6.6E.01 h		1.0E-031 1.0E-0	
	TYVAL			255 255 255 255 255 255 255 255 255 255	1.1E+00 r 3.3E-02 r x x	1.8E+00	8.4E-01 1
	TOXIC	2.0E-03 1.0E+00 3.0E-02		5.7E-08 7.0E-04 1.0E-02 6.0E-02	6.0E-07 6.0E-07 6.0E-07 6.0E-07 6.0E-07	10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2.0E-02 1.5E-03 1.0E-00 1.0E-0
		4,8E-02	1.0E-01	2.5 1.35 4.6 6.02 7.5 6.03	1.1E+000 3.3E-0Z	1,8E+00	

2
1870
2
Smucker
-
80
ry Remediation Gosts:
ŏ
=
옽
#
ĕ
Ē
ž
7
3
늗
둍
5
-
~
E
품
2
A
0
ш

REMEDIAL GOALS (PRGs) Ambient Nr Tap Water (us/m/3) (us/m/3)	2.1E+00 nc 1.8E+01 nc 2.6E+03 nc 7.2E-04 ca 2.5E+03 nc 1.2E-02 ca 1.2E-02 ca 1.2E-02 ca 1.2E-02 ca 1.2E-03 ca 1.2E-04 nc	Color Colo
SES ELIMINARY RE ELIMINARY RE SOL (marks)	2.35	
	3.3E+03 nc 3.0E+03 nc 3.0E+03 nc 3.0E+03 nc 3.1E+03 nc 3.3E+03 nc 3.3E+0	3.3.8.402 nc 3.3.8.403 nc 1.3.403 nc 1.3.403 nc 1.3.403 nc 1.3.8.403 nc
PLANNING PURP TON CHECANCER PRG FICE SATURAL CONTAMINANT Residentia Soil (mark)	Nitrobenzene Nitroterranion Nitrogen dioxide Nitrogen dioxide Nitrogen dioxide Nitrogenerdine 4 Nitrogropane A	Peruthon Pebulate Peruthomodiphenyl ether Peruthomodiphenyl ether Pertachtomodiphenyl ether Pherutomodiphenyl ether Pherutomodiphenyl ether Pherutomodiphenyl ether Phosphine
ST 6=ECAO X=WINDRAWN THROUTE EXTRAPOLATION IN THE STATE OF STATE O	3.7E-04 h 0 0.10 3.4E+00 h 7.0E-02 h 0 0.10 5.8E+00 h 0.10 7.0E-02 h 0.10	2.3E-02; 7.4.0E-03; 7.0 0.10 2.3E-02; 7.4.0E-02; 7.0 0.10 2.3E-03; 7.0 0.10 1.2E-04; 7.3.0E-03; 7.0 0.10 1.3E-04; 7.0 0.10 1.3E-03; 7.0 0.10
Key: I=IR S IN=HEAST TOXIC	1.5E+00 h 1.0E-02 h 1.0E-0	2.3E-02 h 3.0E-03 h 1.9E-03 h 1.9E-03 h 1.9E-04 h 1.9E-0

REWEDIAL GOALS (PRGs) Ambient Ar Tap Water (ug/m/3) (ug/l)	3.7E+01 Inc. 3.7E+02 Inc. 1.8E+03 Inc. 1.5E+03 Inc. 1.5E+03 Inc. 1.5E+03 Inc. 1.5E+03 Inc. 1.5E+02 Inc. 1.5E+04 Inc. 1.5E+
SES MERCELLING LMIT ELIMINARY R Industria Sod (marka)	8. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.
PURPO Self-SOIL SATURATION PR Residentia Soil (marks)	8
PLANINING P	Primiphoe-methy Polybrominated biphenyls Polybrominated biphenyls Polybrominated biphenyls Polybrominated biphenyls Polybrominated biphenyls Polybrominated biphenyls Avchor 1016 Polybridge promatic indicate Berizolelluoranihene Berizolelluoranihene Berizolelluoranihene Berizolelluoranihene Berizolelluoranihene Berizolelluoranihene Perizolelluoranihene Promatika Pr
FOR EXTRAPOLA OIL FACTORS OF SHEET OF S	
Y VAL UES S ISF RED IMMENSE OF MATTER INTERPRETATION OF MATTER INT	7.3E-01 r 7.0E-02 r 7.3E-01 r 7.3E-01 r 7.3E-02 r 7.3E-03 r 7.3E-0
Key: I=IRS h=HEAST e+EC TOXIC)TY oSF oRID II/mg/kg-d) (mg/kg-d) II/i	4.5E+00 h 7.0E-021 7.3E-03 e 7.0E-021 7.5E-03 e 7.0E-031 7.5E-03 e 7.0E-031 7.5E-0

J. Smucker (08/01/94)
w
Goals:
Remediation
Preliminary
×
Region
EPA

	Ambert Ar Tap Water (up/m/3) (up/m/3)	3.3E+02 in 3.3E+03 in 3.3E+03 in 3.3E+02 in 3.3E+02 in 3.3E+03 in
SES	PRELIMINARY REMI	2. 1
PURP(Residential Soi (mg/kg)	1
PLA	CONTAMINANT	Sethonydim Siner and compounds Siner and compounds Sodium azide Sodium metavanadate Sodium metavanadate Sodium metavanadate Sodium metavanadate Sirverie Sir
FOR	OL FACTORS V Oskin VF	
AO x= WITHDRAWK T	ST RID Maked	2.7E-01 r 2.0E-03 r 1.5E-05 r 1.2E-01 r 2.0E-03 r 1.2E-05 r 1.2E-0
10	OSF ORID (Mg/kg-d) (1/1	2.7E-01 h 50E-02 h 1.2E-01 h 2.0E-03 h 1.2E-01 h 2.0E-03 h 1.2E-03

R	GOALS	Amblent Air Tap Water (up/m/3) (up/m/3)	-01 nc 1.1E+00	2.0E-01 cs 2.0E+00 cs 2.3E-01 cs 2.3E-01 cs	1.9E+02	2.0E-01	1.1E+00 ca* 1.6E+00 ca*	2 2	a 6.1E+00	9 5	3.0E+01	3.1E+01	PAC S MEANS	2 2	nc 1.1E+02	nc 1.2E+01	9.7E+00	8 2	3.7E+02	Ca 2.2	1.1E+02 nc	3 35+02 12	7.3E+02 inc		6 1E400 TG 6 4E401 TG	3 7E+04	2.2E-02 ca 2.0E-02 ca	, 100	IC 1.4E+03		7.3E+02 no 1.4E+03 nc	1.1E+04 nc	1.8E+02 Inc 1.8E+03 Inc
SES	ELIMINARY	Industrial Soil (mg/ligi)	2.0E+01 nc	5.6E+01 cs 6.6E+01 cs	5.0E+03 nc	1.5E+00 ca	7.3E+00 Ca*	6.8E+04 Inc	1.7E+02 ca	5 5F+03 nc	8.0E+01 nc	6.4E-03 ca	3F+00 Inc	3.6F+03 net	2.0E+03 Inc	3.5E+01 nc	2.5E+02 Ca*	3 4 1 100	8.8E+03 nc	6.4E+01 Ca	1 25+03 110	1.5E+04 Inc	3.4E+04 nc	3.4E+04 nc	1 7F+04 mg	1.0E+05 Imax	3.5E-03 ca	2.0E+02 nc	9.8E+02 sar	9.8E+02 land	9.8E+02 sat	1.0E+05 max	3.4E+04 Inc
OURPC REFSOIL SATURATION	PR	Residential Soil (mg/kg)	2.0E+00 nc	1.3E+01 ca	5.8E+02 nc	8.6E-01 ca	3.3E+00 cs.	6.5E+03 nc	4.0E+01 CB	5.2E+02 nc	2.3E+01 Inc	2.9E-03 ca		3.6E+03 sel			5.8E+01 C8-	3.3E+00 nc	6.5E+02 nc	4.8E+01 ca	5 4 5 40	6.9E+02 nc	1.5E+03 Inc	1.95+03 170				Š	9.8E+02 860	Ş	Ş	235+04 76	3.3E+03 nc
PL/	CONTAMINANT		Tributyttin code (TBTO)	2.4.6-Inchlorestime hydrochloride	1,2,4-1 richtorobenzene	1,1,2-Trichloroethane	Trichloromethane	2.4.5-Trichlorophenol	2.4.6- inchiorophenol	2-(2,4,5-Trichlorophenoxy) propionic acid	1,1,2-Trichloropropane	17.2.3-Inchiopropare	N	11,1,2-Trichloro-1,2,2-trifluoroethane	맏	Inelhylamine	Trimoffivi mosmosta	1,3,5-Trinitrobenzene	Trinitrophenymethylntramine	2.4.6-Trititotoluene	nager	Vanadium pentoxide	Vanach suitate	Verteum surers	Vindozolin	Vinyl acetate	Vinyl choride	Wanarin	In-Xyrana In-Xyrana	p-Xylene	Xylene (mixed)	Zivo phosphide	Zineb
OR	7	(m ^A 3/kg)	00	0.10		0 6.0E+03			500		0 3.2E+03		0 5.2E+03			3.5E+03		0	0	0 2		į			0	П	0 7.4E+01	_1_	2.45+04		낢		0
HDRAWN FROU	2 2 2 2	(mg/kg-d) C ABS	\rightarrow		i.			-		-		-	5.0E-03 r 1 0.10	1	9	7		0	_	5	-	_		1	_				20E-01 x 11 0.10			90	5.0E-02 r 0 0.1
HEAST GECAO X=VI	<u> </u>	D-Byom/th (D-B		2.96-02 г	-	4.0E-03 5.8E-02	0.05-00	H	1,15-02	03	-03	1.05.00	-03 h	10+	63	t	3.7E-02 F	Н		3.05-02	-03 h	63	196	13	-021		3.0E-01 In	58			004	04	-02
Køy: FIRS h=HEAS		(D-8/16m) D-8/16m)//	3.0E-05	++	+	5.7E-02 4.0E	H	1.0E-0	+	8.0E-03	V NE 100 5.0E	t	5.0E-03	3.0E	3.08	1	3.7E-02/h	Н	1.0E	3.0E-03	7.0E	9.05	200	1.00	2.56		1.9E+00 m	3.KE	2.0€+00		205	3.0E-04	1 5.0E